

Nonexpansive Iterations in Uniformly Convex W -hyperbolic Spaces

Laurențiu Leuştean

ABSTRACT. We propose the class of uniformly convex W -hyperbolic spaces with monotone modulus of uniform convexity (UCW -hyperbolic spaces for short) as an appropriate setting for the study of nonexpansive iterations. UCW -hyperbolic spaces are a natural generalization both of uniformly convex normed spaces and $CAT(0)$ spaces. Furthermore, we apply proof mining techniques to get effective rates of asymptotic regularity for Ishikawa iterations of nonexpansive self-mappings of closed convex subsets in UCW -hyperbolic spaces. These effective results are new even for uniformly convex Banach spaces.

1. Introduction

In this paper we propose the class of uniformly convex W -hyperbolic spaces with monotone modulus of uniform convexity (UCW -hyperbolic spaces for short) as an appropriate setting for the study of nonexpansive iterations. This class of geodesic spaces, which will be defined in Section 2, is a natural generalization both of uniformly convex normed spaces and $CAT(0)$ spaces. As we shall see in Section 2, complete UCW -hyperbolic spaces have very nice properties. Thus, the intersection of any decreasing sequence of bounded closed convex subsets is nonempty (Proposition 2.2) and closed convex subsets are Chebyshev sets (Proposition 2.4).

The asymptotic center technique, introduced by Edelstein [6, 7], is one of the most useful tools in metric fixed point theory of nonexpansive mappings in uniformly convex Banach spaces, due to the fact that bounded sequences have unique asymptotic centers with respect to closed convex subsets. We prove that this basic property is true for complete UCW -hyperbolic spaces, too (Proposition 3.3). The main result of Section 3 is Theorem 3.5, which uses methods involving asymptotic centers to get, for nonexpansive self-mappings $T : C \rightarrow C$ of convex closed subsets of complete UCW -hyperbolic spaces, equivalent characterizations of the fact that T has fixed points in terms of boundedness of different iterations associated

2000 *Mathematics Subject Classification.* Primary 47H10,47H09,03F10.

Key words and phrases. Nonexpansive mappings, metric fixed point theory, asymptotic regularity, uniformly convex hyperbolic spaces, asymptotic centers, proof mining.

with T . As an immediate consequence of Theorem 3.5, we obtain a generalization to complete UCW -hyperbolic spaces of the well-known Browder-Goehde-Kirk Theorem.

In the second part of the paper, we apply proof mining techniques to give effective rates of asymptotic regularity for Ishikawa iterations of nonexpansive self-mappings of closed convex subsets in UCW -hyperbolic spaces. We emphasize that our results are new even for the normed case. By *proof mining* we mean the logical analysis of mathematical proofs with the aim of extracting new numerically relevant information hidden in the proofs. We refer to Kohlenbach's book [16] for details on proof mining.

If $(X, \|\cdot\|)$ is a normed space, $C \subseteq X$ a convex subset of X and $T : C \rightarrow C$ is nonexpansive, then the *Ishikawa iteration* [11] starting with $x \in C$ is defined by

$$(1) \quad x_0 := x, \quad x_{n+1} = (1 - \lambda_n)x_n + \lambda_n T((1 - s_n)x_n + s_n T x_n),$$

where $(\lambda_n), (s_n)$ are sequences in $[0, 1]$. By letting $s_n = 0$ for all $n \in \mathbb{N}$, we get the Krasnoselski-Mann iteration as a special case.

In Section 4, we consider the important problem of asymptotic regularity associated with the Ishikawa iterations:

$$\lim_{n \rightarrow \infty} d(x_n, T x_n) = 0.$$

Our point of departure is the following result, proved by Tan and Xu [27] for uniformly convex Banach spaces and, recently, by Dhompongsa and Panyanak [4] for complete $CAT(0)$ spaces.

PROPOSITION 1.1. *Let X be a uniformly convex Banach space or a complete $CAT(0)$ space, $C \subseteq X$ a bounded closed convex subset and $T : C \rightarrow C$ be nonexpansive. Assume that $\sum_{n=0}^{\infty} \lambda_n(1 - \lambda_n)$ diverges, $\limsup_n s_n < 1$ and $\sum_{n=0}^{\infty} s_n(1 - \lambda_n)$ converges.*

Then for all $x \in C$,

$$\lim_{n \rightarrow \infty} d(x_n, T x_n) = 0.$$

Using proof mining methods we obtain a quantitative version (Theorem 4.7) of a two-fold generalization of the above proposition:

- firstly, we consider UCW -hyperbolic spaces;
- secondly, we assume that T has a fixed point instead of assuming the boundedness of C .

The idea is to combine methods used in [20] to obtain effective rates of asymptotic regularity for Krasnoselski-Mann iterates with the ones used in [21] to get rates of asymptotic regularity for Halpern iterates.

In this way, we provide for the first time (even for the normed case) effective rates of asymptotic regularity for the Ishikawa iterates, that is, rates of convergence of $(d(x_n, T x_n))$ towards 0.

For bounded C (Corollary 4.9), the rate of asymptotic regularity is uniform in the nonexpansive mapping T and the starting point $x \in C$ of the iteration, and it depends on C only via its diameter and on the space X only via a monotone modulus of uniform convexity.

2. UCW-hyperbolic spaces

We work in the setting of hyperbolic spaces as introduced by Kohlenbach [15]. In order to distinguish them from Gromov hyperbolic spaces [2] or from other notions of 'hyperbolic space' which can be found in the literature (see for example [13, 8, 24]), we shall call them W -hyperbolic spaces.

A W -hyperbolic space (X, d, W) is a metric space (X, d) together with a convexity mapping $W : X \times X \times [0, 1] \rightarrow X$ satisfying

$$\begin{aligned} (W1) \quad & d(z, W(x, y, \lambda)) \leq (1 - \lambda)d(z, x) + \lambda d(z, y), \\ (W2) \quad & d(W(x, y, \lambda), W(x, y, \tilde{\lambda})) = |\lambda - \tilde{\lambda}| \cdot d(x, y), \\ (W3) \quad & W(x, y, \lambda) = W(y, x, 1 - \lambda), \\ (W4) \quad & d(W(x, z, \lambda), W(y, w, \lambda)) \leq (1 - \lambda)d(x, y) + \lambda d(z, w). \end{aligned}$$

The convexity mapping W was first considered by Takahashi in [26], where a triple (X, d, W) satisfying (W1) is called a *convex metric space*. If (X, d, W) satisfies (W1)–(W3), then we get the notion of *space of hyperbolic type* in the sense of Goebel and Kirk [8]. (W4) was already considered by Itoh [12] under the name 'condition III' and it is used by Reich and Shafrir [24] and Kirk [13] to define their notions of hyperbolic space. We refer to [16, pp. 384-387] for a detailed discussion.

The class of W -hyperbolic spaces includes normed spaces and convex subsets thereof, the Hilbert ball (see [9] for a book treatment) as well as CAT(0) spaces.

If $x, y \in X$ and $\lambda \in [0, 1]$, then we use the notation $(1 - \lambda)x \oplus \lambda y$ for $W(x, y, \lambda)$. The following holds even for the more general setting of convex metric spaces [26]: for all $x, y \in X$ and $\lambda \in [0, 1]$,

$$(2) \quad d(x, (1 - \lambda)x \oplus \lambda y) = \lambda d(x, y), \text{ and } d(y, (1 - \lambda)x \oplus \lambda y) = (1 - \lambda)d(x, y).$$

As a consequence, $1x \oplus 0y = x$, $0x \oplus 1y = y$ and $(1 - \lambda)x \oplus \lambda x = \lambda x \oplus (1 - \lambda)x = x$.

For all $x, y \in X$, we shall denote by $[x, y]$ the set $\{(1 - \lambda)x \oplus \lambda y : \lambda \in [0, 1]\}$. Thus, $[x, x] = \{x\}$ and for $x \neq y$, the mapping

$$\gamma_{xy} : [0, d(x, y)] \rightarrow \mathbb{R}, \quad \gamma(\alpha) = \left(1 - \frac{\alpha}{d(x, y)}\right)x \oplus \frac{\alpha}{d(x, y)}y$$

is a geodesic satisfying $\gamma_{xy}([0, d(x, y)]) = [x, y]$. That is, any W -hyperbolic space is a geodesic space.

A subset $C \subseteq X$ is *convex* if $[x, y] \subseteq C$ for all $x, y \in C$. A nice feature of our setting is that any convex subset is itself a W -hyperbolic space with the restriction of d and W to C . It is easy to see that open and closed balls are convex. Moreover, using (W4), we get that the closure of a convex subset of a W -hyperbolic space is again convex.

If C is a convex subset of X , then a function $f : C \rightarrow \mathbb{R}$ is said to be *convex* if

$$f((1 - \lambda)x \oplus \lambda y) \leq (1 - \lambda)f(x) + \lambda f(y)$$

for all $x, y \in C, \lambda \in [0, 1]$.

One of the most important classes of Banach spaces are the uniformly convex ones, introduced by Clarkson in the 30's [3]. Following [9, p. 105], we can define uniform convexity for W -hyperbolic spaces, too.

A W -hyperbolic space (X, d, W) is *uniformly convex* [20] if for any $r > 0$ and any $\varepsilon \in (0, 2]$ there exists $\delta \in (0, 1]$ such that for all $a, x, y \in X$,

$$(3) \quad \left. \begin{array}{l} d(x, a) \leq r \\ d(y, a) \leq r \\ d(x, y) \geq \varepsilon r \end{array} \right\} \Rightarrow d\left(\frac{1}{2}x \oplus \frac{1}{2}y, a\right) \leq (1 - \delta)r.$$

A mapping $\eta : (0, \infty) \times (0, 2] \rightarrow (0, 1]$ providing such a $\delta := \eta(r, \varepsilon)$ for given $r > 0$ and $\varepsilon \in (0, 2]$ is called a *modulus of uniform convexity*. We call η *monotone* if it decreases with r (for a fixed ε).

LEMMA 2.1. [20, 18] *Let (X, d, W) be a uniformly convex W -hyperbolic space and η be a modulus of uniform convexity. Assume that $r > 0, \varepsilon \in (0, 2], a, x, y \in X$ are such that*

$$d(x, a) \leq r, \quad d(y, a) \leq r \quad \text{and} \quad d(x, y) \geq \varepsilon r.$$

Then for any $\lambda \in [0, 1]$,

- (i) $d((1 - \lambda)x \oplus \lambda y, a) \leq (1 - 2\lambda(1 - \lambda)\eta(r, \varepsilon))r$;
- (ii) for any $\psi \in (0, 2]$ such that $\psi \leq \varepsilon$,

$$d((1 - \lambda)x \oplus \lambda y, a) \leq (1 - 2\lambda(1 - \lambda)\eta(r, \psi))r$$

- (iii) for any $s \geq r$,

$$d((1 - \lambda)x \oplus \lambda y, a) \leq \left(1 - 2\lambda(1 - \lambda)\eta\left(s, \frac{\varepsilon r}{s}\right)\right)s$$

- (iv) if η is monotone, then for any $s \geq r$,

$$d((1 - \lambda)x \oplus \lambda y, a) \leq (1 - 2\lambda(1 - \lambda)\eta(s, \varepsilon))r.$$

PROOF. (i) is a generalization to our setting of a result due to Groetsch [10]. We refer to [20, Lemma 7] for the proof.

(ii), (iii) are immediate; see [18, Lemma 2.1].

(iv) Use (i) and the fact that $\eta(r, \varepsilon) \geq \eta(s, \varepsilon)$, hence $1 - 2\lambda(1 - \lambda)\eta(r, \varepsilon) \leq 1 - 2\lambda(1 - \lambda)\eta(s, \varepsilon)$. \square

We shall refer to uniformly convex W -hyperbolic spaces with a monotone modulus of uniform convexity as *UCW-hyperbolic spaces*. It turns out [20] that CAT(0) spaces are UCW-hyperbolic spaces with modulus of uniform convexity $\eta(r, \varepsilon) = \varepsilon^2/8$ quadratic in ε . Thus, UCW-hyperbolic spaces are a natural generalization of both uniformly convex normed spaces and CAT(0) spaces.

For the rest of this section, (X, d, W) is a complete UCW-hyperbolic space and η is a monotone modulus of uniform convexity.

PROPOSITION 2.2. [18, Proposition 2.2] *The intersection of any decreasing sequence of nonempty bounded closed convex subsets of X is nonempty.*

The next proposition, inspired by [9, Proposition 2.2], is essential for what follows.

PROPOSITION 2.3. *Let C be a closed convex subset of X , $f : C \rightarrow [0, \infty)$ be convex and lower semicontinuous. Assume moreover that for all sequences (x_n) in C ,*

$$\lim_{n \rightarrow \infty} d(x_n, a) = \infty \quad \text{for some } a \in X \quad \text{implies} \quad \lim_{n \rightarrow \infty} f(x_n) = \infty.$$

Then f attains its minimum on C . If, in addition,

$$f\left(\frac{1}{2}x \oplus \frac{1}{2}y\right) < \max\{f(x), f(y)\}$$

for all $x \neq y$, then f attains its minimum at exactly one point.

PROOF. Let α be the infimum of f on C and define

$$C_n := \left\{x \in C \mid f(x) \leq \alpha + \frac{1}{n}\right\}$$

for all $n \in \mathbb{N}$. It is easy to see that we can apply Proposition 2.2 to the sequence $(C_n)_{n \in \mathbb{N}}$ to get the existence of $x^* \in \bigcap_{n \in \mathbb{N}} C_n$. It follows that $f(x^*) \leq \alpha + \frac{1}{n}$ for all $n \geq 1$, hence $f(x^*) \leq \alpha$. Since α is the infimum of f , we can conclude that $f(x^*) = \alpha$, that is, f attains its minimum on C . The second part of the conclusion is immediate. If f attains its minimum at two points $x^* \neq y^*$, then $\frac{1}{2}x^* \oplus \frac{1}{2}y^* \in C$, since C is convex, but $f\left(\frac{1}{2}x^* \oplus \frac{1}{2}y^*\right) < \max\{f(x^*), f(y^*)\} = \alpha$, which is a contradiction. \square

Let us recall that a subset C of a metric space (X, d) is called a *Chebyshev set* if to each point $x \in X$ there corresponds a unique point $z \in C$ such that $d(x, z) = d(x, C) (= \inf\{d(x, y) \mid y \in C\})$. If C is a Chebyshev set, one can define the *nearest point projection* $P : X \rightarrow C$ by assigning z to x .

PROPOSITION 2.4. *Every closed convex subset C of X is a Chebyshev set.*

PROOF. Let $x \in X$ and define $f : C \rightarrow [0, \infty)$, $f(y) = d(x, y)$. Then f is continuous, convex (by (W1)), and for any sequence (y_n) in C , $\lim_{n \rightarrow \infty} d(y_n, a) = \infty$ for some $a \in X$ implies $\lim_{n \rightarrow \infty} f(y_n) = \infty$, since $f(y_n) = d(x, y_n) \geq d(y_n, a) - d(x, a)$. Moreover, let $y \neq z \in C$ and denote $M := \max\{f(y), f(z)\} > 0$. Then

$$d(x, y), d(x, z) \leq M \quad \text{and} \quad d(y, z) \geq \varepsilon \cdot M,$$

where $\varepsilon := \frac{d(y, z)}{M}$ and $0 < \varepsilon \leq \frac{d(x, y) + d(x, z)}{M} \leq 2$. Hence, by uniform convexity it follows that

$$d\left(\frac{1}{2}y \oplus \frac{1}{2}z, x\right) \leq (1 - \eta(M, \varepsilon)) \cdot M < M.$$

Thus, f satisfies all the hypotheses of Proposition 2.3, so we can apply it to conclude that f has a unique minimum. Hence, C is a Chebyshev set. \square

3. Asymptotic centers and fixed point theory of nonexpansive mappings

In the sequel, we recall basic facts about asymptotic centers. We refer to [6, 7, 9] for all the unproved results.

Let (X, d) be a metric space, (x_n) be a bounded sequence in X and $C \subseteq X$ be a subset of X . We define the following functionals:

$$\begin{aligned} r_m(\cdot, (x_n)) : X &\rightarrow [0, \infty), & r_m(y, (x_n)) &= \sup\{d(y, x_n) \mid n \geq m\} \\ & & &\text{for } m \in \mathbb{N}, \\ r(\cdot, (x_n)) : X &\rightarrow [0, \infty), & r(y, (x_n)) &= \limsup_n d(y, x_n) = \inf_m r_m(y, (x_n)) \\ & & &= \lim_{m \rightarrow \infty} r_m(y, (x_n)). \end{aligned}$$

The following lemma collects some basic properties of the above functionals.

LEMMA 3.1. *Let $y \in X$.*

- (i) $r_m(\cdot, (x_n))$ is nonexpansive for all $m \in \mathbb{N}$;
- (ii) $r(\cdot, (x_n))$ is continuous and $r(y, (x_n)) \rightarrow \infty$ whenever $d(y, a) \rightarrow \infty$ for some $a \in X$;
- (iii) $r(y, (x_n)) = 0$ if and only if $\lim_{n \rightarrow \infty} x_n = y$;
- (iv) if (X, d, W) is a convex metric space and C is convex, then $r(\cdot, (x_n))$ is a convex function.

The asymptotic radius of (x_n) with respect to C is defined by

$$r(C, (x_n)) = \inf\{r(y, (x_n)) \mid y \in C\}.$$

The asymptotic radius of (x_n) , denoted by $r((x_n))$, is the asymptotic radius of (x_n) with respect to X , that is, $r((x_n)) = r(X, (x_n))$.

A point $c \in C$ is said to be an asymptotic center of (x_n) with respect to C if

$$r(c, (x_n)) = r(C, (x_n)) = \min\{r(y, (x_n)) \mid y \in C\}.$$

We denote with $A(C, (x_n))$ the set of asymptotic centers of (x_n) with respect to C . When $C = X$, we call c an asymptotic center of (x_n) and we use the notation $A((x_n))$ for $A(X, (x_n))$.

The following lemma, inspired by [7, Theorem 1], turns out to be very useful in the following.

LEMMA 3.2. *Let (x_n) be a bounded sequence in X with $A(C, (x_n)) = \{c\}$ and $(\alpha_n), (\beta_n)$ be real sequences such that $\alpha_n \geq 0$ for all $n \in \mathbb{N}$, $\limsup_n \alpha_n \leq 1$ and $\limsup_n \beta_n \leq 0$.*

Assume that $y \in C$ is such that there exist $p, N \in \mathbb{N}$ satisfying

$$\forall n \geq N \left(d(y, x_{n+p}) \leq \alpha_n d(c, x_n) + \beta_n \right).$$

Then $y = c$.

PROOF. We have that

$$\begin{aligned} r(y, (x_n)) &= \limsup_n d(y, x_n) = \limsup_n d(y, x_{n+p}) \leq \limsup_n (\alpha_n d(c, x_n) + \beta_n) \\ &\leq \limsup_n \alpha_n \cdot \limsup_n d(c, x_n) + \limsup_n \beta_n \leq \limsup_n d(c, x_n) \\ &= r(c, (x_n)). \end{aligned}$$

Since c is unique with the property that $r(c, (x_n)) = \min\{r(z, (x_n)) \mid z \in C\}$, we must have $y = c$. \square

In general, the set $A(C, (x_n))$ of asymptotic centers of a bounded sequence (x_n) with respect to $C \subseteq X$ may be empty or even contain infinitely many points.

The following result shows that in the case of complete UCW-hyperbolic spaces, the situation is as nice as for uniformly convex Banach spaces (see, for example, [9, Theorem 4.1]).

PROPOSITION 3.3. *Let (X, d, W) be a complete UCW-hyperbolic space. Every bounded sequence (x_n) in X has a unique asymptotic center with respect to any closed convex subset C of X .*

PROOF. Let η be a monotone modulus of uniform convexity. We apply Proposition 2.3 to show that the function $r(\cdot, (x_n)) : C \rightarrow [0, \infty)$ attains its minimum at exactly one point. By Lemma 3.1, it remains to prove that

$$r\left(\frac{1}{2}y \oplus \frac{1}{2}z, (x_n)\right) < \max\{r(y, (x_n)), r(z, (x_n))\} \quad \text{whenever } y, z \in C, y \neq z.$$

Let $M := \max\{r(y, (x_n)), r(z, (x_n))\} > 0$. For every $\varepsilon \in (0, 1]$ there exists N such that $d(y, x_n), d(z, x_n) \leq M + \varepsilon \leq M + 1$ for all $n \geq N$. Moreover, $d(y, z) = \frac{d(y, z)}{M + \varepsilon} \cdot (M + \varepsilon) \geq \frac{d(y, z)}{M + 1} \cdot (M + \varepsilon)$. Thus, we can apply Lemma 2.1. (iv) to get that for all $n \geq N$,

$$d\left(\frac{1}{2}y \oplus \frac{1}{2}z, x_n\right) \leq \left(1 - \eta\left(M + 1, \frac{d(y, z)}{M + 1}\right)\right)(M + \varepsilon),$$

hence

$$r\left(\frac{1}{2}y \oplus \frac{1}{2}z, (x_n)\right) \leq \left(1 - \eta\left(M + 1, \frac{d(y, z)}{M + 1}\right)\right)(M + \varepsilon).$$

By letting $\varepsilon \rightarrow 0$, it follows that

$$r\left(\frac{1}{2}y \oplus \frac{1}{2}z, (x_n)\right) \leq \left(1 - \eta\left(M + 1, \frac{d(y, z)}{M + 1}\right)\right) \cdot M < M.$$

This completes the proof. □

In the sequel, we assume that (X, d, W) is a W -hyperbolic space, $C \subseteq X$ is convex and $T : C \rightarrow C$ is *nonexpansive*, that is,

$$d(Tx, Ty) \leq d(x, y)$$

for all $x, y \in C$. We shall denote with $\text{Fix}(T)$ the set of fixed points of T .

For any $\lambda \in (0, 1]$, the *averaged mapping* T_λ is defined by

$$T_\lambda : C \rightarrow C, \quad T_\lambda(x) = (1 - \lambda)x \oplus \lambda Tx.$$

It is easy to see that T_λ is also nonexpansive and that $\text{Fix}(T) = \text{Fix}(T_\lambda)$.

The *Krasnoselski iteration* [19, 25] (x_n) starting with $x \in C$ is defined as the Picard iteration $(T_\lambda^n(x))$ of T_λ , that is,

$$(4) \quad x_0 := x, \quad x_{n+1} := (1 - \lambda)x_n \oplus \lambda Tx_n.$$

By allowing general sequences (λ_n) in $[0, 1]$, we get the *Krasnoselski-Mann iteration* [22] (called *segmenting Mann iterate* in [10]) (x_n) starting with $x \in C$:

$$(5) \quad x_0 := x, \quad x_{n+1} := (1 - \lambda_n)x_n \oplus \lambda_n Tx_n.$$

We refer to [1, 14, 17, 24] for results on the general behavior of the Krasnoselski-Mann iterations.

The following lemma collects some known properties of Krasnoselski-Mann iterates in W -hyperbolic spaces. For the sake of completeness we prove them here.

LEMMA 3.4. *Assume that $(x_n), (y_n)$ are the Krasnoselski-Mann iterations starting with $x, y \in C$. Then*

- (i) $(d(x_n, y_n))$ is decreasing;
- (ii) if p is a fixed point of T , then $(d(x_n, p))$ is decreasing;
- (iii) $d(x_{n+1}, Ty) \leq d(x_n, y) + (1 - \lambda_n)d(y, Ty)$ for all $n \in \mathbb{N}$.

PROOF.

$$\begin{aligned}
d(x_{n+1}, y_{n+1}) &\leq (1 - \lambda_n)d(x_n, y_n) + \lambda_n d(Tx_n, Ty_n) \quad \text{by (W4)} \\
&\leq d(x_n, y_n), \quad \text{since } T \text{ is nonexpansive,} \\
d(x_{n+1}, p) &\leq (1 - \lambda_n)d(x_n, p) + \lambda_n d(Tx_n, p) \quad \text{by (W1)} \\
&= (1 - \lambda_n)d(x_n, p) + \lambda_n d(Tx_n, Tp) \\
&\leq (1 - \lambda_n)d(x_n, p) + \lambda_n d(x_n, p) = d(x_n, p), \\
d(x_{n+1}, Ty) &\leq (1 - \lambda_n)d(x_n, Ty) + \lambda_n d(Tx_n, Ty) \quad \text{by (W1)} \\
&\leq (1 - \lambda_n)d(x_n, y) + (1 - \lambda_n)d(Ty, y) + \lambda_n d(x_n, y) \\
&\leq d(x_n, y) + (1 - \lambda_n)d(Ty, y).
\end{aligned}$$

□

We can prove now the main theorem of this section.

THEOREM 3.5. *Let (X, d, W) be a complete UCW-hyperbolic space, $C \subseteq X$ a convex closed subset and $T : C \rightarrow C$ be nonexpansive.*

The following are equivalent

- (i) T has a fixed point;
- (ii) there exists a bounded sequence (u_n) in C such that $\lim_{n \rightarrow \infty} d(u_n, Tu_n) = 0$;
- (iii) the sequence $(T^n x)$ of Picard iterates is bounded for some $x \in C$;
- (iv) the sequence $(T^n x)$ of Picard iterates is bounded for all $x \in C$;
- (v) the Krasnoselski-Mann iteration (x_n) is bounded for some $x \in C$ and for (λ_n) in $[0, 1]$ satisfying one of the following conditions:
 - (a) $\lambda_n = \lambda \in (0, 1]$;
 - (b) $\lim_{n \rightarrow \infty} \lambda_n = 1$;
 - (c) $\limsup_n \lambda_n < 1$ and $\sum_{n=0}^{\infty} \lambda_n$ diverges;
- (vi) the Krasnoselski-Mann iteration (x_n) is bounded for all $x \in C$ and all (λ_n) in $[0, 1]$.

PROOF. (i) \Rightarrow (ii) Let p be a fixed point of T and define $u_n := p$ for all $n \in \mathbb{N}$.

(ii) \Rightarrow (i) By Proposition 3.3, (u_n) has a unique asymptotic center c with respect to C . We get that for all $n \in \mathbb{N}$,

$$d(Tc, u_n) \leq d(Tc, Tu_n) + d(Tu_n, u_n) \leq d(c, u_n) + d(Tu_n, u_n).$$

We can apply now Lemma 3.2 with $y := Tc$ and $p := N := 0$, $\alpha_n := 1, \beta_n := d(u_n, Tu_n)$ to get that $Tc = c$.

(i) \Rightarrow (iii) If p is a fixed point of T , then $T^n p = p$ for all $n \in \mathbb{N}$.

(iii) \Rightarrow (iv) Apply the fact that, since T is nonexpansive, $d(T^n x, T^n y) \leq d(x, y)$ for all $x, y \in C$.

(iv) \Rightarrow (i) Let $c \in C$ be the unique asymptotic center of $(T^n x)$. Then for all $n \in \mathbb{N}$,

$$d(Tc, T^{n+1}x) \leq d(c, T^n x),$$

hence we can apply Lemma 3.2 with $y := Tc, x_n := T^n x$ and $p := 1, N := 0, \alpha_n := 1, \beta_n := 0$ to get that $Tc = c$.

(i) \Rightarrow (vi) Let p be a fixed point of T . Then for any $x \in C, (\lambda_n)$ in $[0, 1]$, the sequence $(d(x_n, p))$ is decreasing, hence bounded from above by $d(x, p)$.

(vi) \Rightarrow (v) Obvious.

- (v) \Rightarrow (i)
- (a) If $\lambda_n = \lambda \in (0, 1]$, then (x_n) is the Krasnoselski iteration, hence the Picard iteration $T_\lambda^n(x)$ of the nonexpansive mapping T_λ . Apply now (iii) \Rightarrow (i) and the fact that $\text{Fix}(T) = \text{Fix}(T_\lambda)$ to get that T has fixed points.
- (b) Assume now that $\lim_{n \rightarrow \infty} \lambda_n = 1$ and let $c \in C$ be the asymptotic center of (x_n) . By Lemma 3.4.(iii), we get that

$$d(Tc, x_{n+1}) \leq d(c, x_n) + (1 - \lambda_n)d(c, Tc).$$

Apply now Lemma 3.2 with $y := Tc$ and $p := 1, N := 0, \alpha_n := 1, \beta_n := (1 - \lambda_n)d(c, Tc)$ to get that $Tc = c$.

- (c) Finally, if (λ_n) is bounded away from 1 and divergent in sum, then $\lim d(x_n, Tx_n) = 0$ by [17, Theorem 3.21], proved even for W -hyperbolic space. Hence (ii) holds. □

As an immediate consequence we obtain the generalization to complete UCW -hyperbolic spaces of the well-known Browder-Goehde-Kirk Theorem.

COROLLARY 3.6. *Let (X, d, W) be a complete UCW -hyperbolic space, $C \subseteq X$ a bounded convex closed subset and $T : C \rightarrow C$ be nonexpansive. Then T has fixed points.*

Theorem 3.5 states that, under some hypotheses on (λ_n) , if the Krasnoselski-Mann iteration starting from some point $x \in C$ is bounded, then T has a fixed point. For Banach spaces, a far-reaching result of this type was obtained by Reich [23] for the very general Mann iterates [22], extending work of Dotson and Mann [5] done in the setting of uniformly convex Banach spaces.

4. Rates of asymptotic regularity for the Ishikawa iterations

Let (X, d, W) be a W -hyperbolic space, $C \subseteq X$ a convex subset of X and $T : C \rightarrow C$ nonexpansive.

As in the case of normed spaces, we can define the *Ishikawa iteration* [11] starting with $x \in C$ by

$$(6) \quad x_0 := x, \quad x_{n+1} = (1 - \lambda_n)x_n \oplus \lambda_n T((1 - s_n)x_n \oplus s_n Tx_n),$$

where $(\lambda_n), (s_n)$ are sequences in $[0, 1]$. By letting $s_n = 0$ for all $n \in \mathbb{N}$, we get the Krasnoselski-Mann iteration as a special case.

We shall use the following notations

$$y_n := (1 - s_n)x_n \oplus s_n Tx_n$$

and

$$T_n : C \rightarrow C, \quad T_n(x) = (1 - \lambda_n)x \oplus \lambda_n T((1 - s_n)x \oplus s_n Tx).$$

Then

$$x_{n+1} = (1 - \lambda_n)x_n \oplus \lambda_n Ty_n = T_n x_n,$$

and it is easy to see that $\text{Fix}(T) \subseteq \text{Fix}(T_n)$ for all $n \in \mathbb{N}$.

Before proving the main technical lemma, we give some basic properties of Ishikawa iterates, which hold even in the very general setting of W -hyperbolic spaces. Their proofs follow closely the ones of the corresponding properties in

uniformly convex Banach spaces [27] or CAT(0) spaces [4], but, for the sake of completeness, we include the details.

LEMMA 4.1.

(i) For all $n \in \mathbb{N}$,

$$(7) \quad d(x_n, x_{n+1}) = \lambda_n d(x_n, Ty_n), \quad d(Ty_n, x_{n+1}) = (1 - \lambda_n) d(x_n, Ty_n),$$

$$(8) \quad d(y_n, x_n) = s_n d(x_n, Tx_n), \quad d(y_n, Tx_n) = (1 - s_n) d(x_n, Tx_n),$$

$$(9) \quad (1 - s_n) d(x_n, Tx_n) \leq d(x_n, Ty_n) \leq (1 + s_n) d(x_n, Tx_n),$$

$$(10) \quad d(y_n, Ty_n) \leq d(x_n, Tx_n),$$

$$(11) \quad d(x_{n+1}, Tx_{n+1}) \leq (1 + 2s_n(1 - \lambda_n)) d(x_n, Tx_n).$$

(ii) T_n is nonexpansive for all $n \in \mathbb{N}$;

(iii) For all $p \in \text{Fix}(T)$, the sequence $(d(x_n, p))$ is decreasing, for all $n \in \mathbb{N}$,

$$d(y_n, p) \leq d(x_n, p) \quad \text{and} \quad d(x_n, Ty_n), d(x_n, Tx_n) \leq 2d(x_n, p).$$

PROOF. (i) (7) and (8) follow from (2).

$$\begin{aligned} d(x_n, Tx_n) &\leq d(x_n, Ty_n) + d(Ty_n, Tx_n) \leq d(x_n, Ty_n) + d(x_n, y_n) \\ &= d(x_n, Ty_n) + s_n d(x_n, Tx_n) \quad \text{by (8)}, \end{aligned}$$

hence $(1 - s_n) d(x_n, Tx_n) \leq d(x_n, Ty_n)$.

$$\begin{aligned} d(x_n, Ty_n) &\leq d(x_n, Tx_n) + d(Tx_n, Ty_n) \leq d(x_n, Tx_n) + d(x_n, y_n) \\ &= (1 + s_n) d(x_n, Tx_n) \quad \text{by (8)}. \end{aligned}$$

$$\begin{aligned} d(y_n, Ty_n) &\leq (1 - s_n) d(x_n, Ty_n) + s_n d(Tx_n, Ty_n) \quad \text{by (W1)} \\ &\leq (1 - s_n)(1 + s_n) d(x_n, Tx_n) + s_n d(x_n, y_n) \quad \text{by (9)} \\ &= d(x_n, Tx_n) \quad \text{by (8)}. \end{aligned}$$

Let us prove now (11). First, let us remark that

$$\begin{aligned} d(x_n, Tx_{n+1}) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, Tx_{n+1}) \\ &= \lambda_n d(x_n, Ty_n) + d(x_{n+1}, Tx_{n+1}) \quad \text{by (7)} \end{aligned}$$

and

$$d(y_n, x_{n+1}) \leq (1 - \lambda_n) d(y_n, x_n) + \lambda_n d(y_n, Ty_n) \quad \text{by (W1)}.$$

Moreover,

$$\begin{aligned} d(x_{n+1}, Tx_{n+1}) &\leq (1 - \lambda_n) d(x_n, Tx_{n+1}) + \lambda_n d(Ty_n, Tx_{n+1}) \quad \text{by (W1)} \\ &\leq (1 - \lambda_n) [d(x_n, x_{n+1}) + d(x_{n+1}, Tx_{n+1})] + \lambda_n d(y_n, x_{n+1}) \\ &\leq (1 - \lambda_n) d(x_{n+1}, Tx_{n+1}) + (1 - \lambda_n) \lambda_n d(x_n, Ty_n) + \\ &\quad + \lambda_n (1 - \lambda_n) d(y_n, x_n) + \lambda_n^2 d(y_n, Ty_n) \\ &\quad \text{by (7) and (W1);} \end{aligned}$$

hence

$$\begin{aligned} d(x_{n+1}, Tx_{n+1}) &\leq (1 - \lambda_n) d(x_n, Ty_n) + (1 - \lambda_n) d(y_n, x_n) + \lambda_n d(y_n, Ty_n) \\ &\leq (1 - \lambda_n)(1 + s_n) d(x_n, Tx_n) + (1 - \lambda_n) s_n d(x_n, Tx_n) \\ &\quad + \lambda_n d(x_n, Tx_n) \quad \text{by (9), (8) and (10)} \\ &= (1 + 2s_n(1 - \lambda_n)) d(x_n, Tx_n). \end{aligned}$$

(ii)

$$\begin{aligned} d(T_n x, T_n y) &\leq (1 - \lambda_n)d(x, y) + \lambda_n d(T((1 - s_n)x \oplus s_n T x), T((1 - s_n)y \oplus s_n T y)) \\ &\leq (1 - \lambda_n)d(x, y) + \lambda_n [(1 - s_n)d(x, y) + s_n d(T x, T y)] \\ &\quad \text{by (W4)} \\ &\leq (1 - \lambda_n)d(x, y) + \lambda_n [(1 - s_n)d(x, y) + s_n d(x, y)] = d(x, y). \end{aligned}$$

(iii)

$$\begin{aligned} d(x_{n+1}, p) &= d(T_n x_n, T_n p) \leq d(x_n, p), \\ d(y_n, p) &\leq (1 - s_n)d(x_n, p) + s_n d(T x_n, p) = (1 - s_n)d(x_n, p) + s_n d(T x_n, T p) \\ &\leq d(x_n, p), \\ d(x_n, T x_n) &\leq d(x_n, p) + d(T x_n, p) \leq 2d(x_n, p), \\ d(x_n, T y_n) &\leq d(x_n, p) + d(T y_n, p) \leq d(x_n, p) + d(y_n, p) \leq 2d(x_n, p). \end{aligned}$$

□

LEMMA 4.2. (**Main technical lemma**) *Assume that (X, d, W) is a UCW-hyperbolic space with a monotone modulus of uniform convexity η and $p \in \text{Fix}(T)$. Let $x \in C, n \in \mathbb{N}$.*

(i) *If $\gamma, \beta, \tilde{\beta}, a > 0$ are such that*

$$\gamma \leq d(x_n, p) \leq \beta, \tilde{\beta} \quad \text{and} \quad a \leq d(x_n, T y_n),$$

then

$$d(x_{n+1}, p) \leq d(x_n, p) - 2\gamma\lambda_n(1 - \lambda_n)\eta\left(\tilde{\beta}, \frac{a}{\beta}\right).$$

(ii) *Assume moreover that η can be written as $\eta(r, \varepsilon) = \varepsilon \cdot \tilde{\eta}(r, \varepsilon)$ such that $\tilde{\eta}$ increases with ε (for a fixed r). If $\delta, a > 0$ are such that*

$$d(x_n, p) \leq \delta \quad \text{and} \quad a \leq d(x_n, T y_n),$$

then

$$d(x_{n+1}, p) \leq d(x_n, p) - 2a\lambda_n(1 - \lambda_n)\tilde{\eta}\left(\delta, \frac{a}{\delta}\right).$$

PROOF. (i) First, let us remark that, using Lemma 4.1.(iii),

$$\begin{aligned} d(T y_n, p) &= d(T y_n, T p) \leq d(y_n, p) \leq d(x_n, p) \leq \beta, \tilde{\beta}, \\ d(x_n, T y_n) \geq a &= \left(\frac{a}{\beta}\right) \cdot \beta \geq \left(\frac{a}{\beta}\right) \cdot d(x_n, p), \quad \text{and} \\ 0 < a \leq d(x_n, T y_n) &\leq 2d(x_n, p) \leq 2\beta, \quad \text{so } \frac{a}{\beta} \in (0, 2]. \end{aligned}$$

Thus, we can apply Lemma 2.1.(iv) with $r := d(x_n, p), s := \tilde{\beta}, \varepsilon := \frac{a}{\tilde{\beta}}$ to obtain

$$\begin{aligned} d(x_{n+1}, p) &= d((1 - \lambda_n)x_n \oplus \lambda_n T y_n, p) \\ &\leq \left(1 - 2\lambda_n(1 - \lambda_n)\eta\left(\tilde{\beta}, \frac{a}{\beta}\right)\right) d(x_n, p) \\ &= d(x_n, p) - 2\lambda_n(1 - \lambda_n)\eta\left(\tilde{\beta}, \frac{a}{\beta}\right) d(x_n, p) \\ &\leq d(x_n, p) - 2\gamma\lambda_n(1 - \lambda_n)\eta\left(\tilde{\beta}, \frac{a}{\beta}\right), \end{aligned}$$

since $d(x_n, p) \geq \gamma$ by hypothesis.

(ii) Since, by Lemma 4.1.(iii), $0 < a \leq d(x_n, Ty_n) \leq 2d(x_n, p)$, we can apply (i) with $\gamma := \beta := d(x_n, p) > 0$ and $\tilde{\beta} := \delta$ to get that

$$\begin{aligned} d(x_{n+1}, p) &\leq d(x_n, p) - 2d(x_n, p)\lambda_n(1 - \lambda_n)\eta\left(\delta, \frac{a}{d(x_n, p)}\right) \\ &= d(x_n, p) - 2a\lambda_n(1 - \lambda_n)\tilde{\eta}\left(\delta, \frac{a}{d(x_n, p)}\right) \\ &\leq d(x_n, p) - 2a\lambda_n(1 - \lambda_n)\tilde{\eta}\left(\delta, \frac{a}{\delta}\right), \end{aligned}$$

since $\frac{a}{\delta} \leq \frac{a}{d(x_n, p)}$ and $\tilde{\eta}$ increases with ε by hypothesis. \square

We recall some terminology. Let $(a_n)_{n \geq 0}$ be a sequence of real numbers. A *rate of divergence* of a divergent series $\sum_{n=0}^{\infty} a_n$ is a function $\theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\sum_{i=0}^{\theta(n)} a_i \geq n$ for all $n \in \mathbb{N}$.

If $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$, then a function $\gamma : (0, \infty) \rightarrow \mathbb{N}$ is called

- a *Cauchy modulus* of (a_n) if $|a_{\gamma(\varepsilon)+n} - a_{\gamma(\varepsilon)}| < \varepsilon$ for all $\varepsilon > 0, n \in \mathbb{N}$;
- a *rate of convergence* of (a_n) if $|a_{\gamma(\varepsilon)+n} - a| < \varepsilon$ for all $\varepsilon > 0, n \in \mathbb{N}$.

A *Cauchy modulus* of a convergent series $\sum_{n=0}^{\infty} a_n$ is a Cauchy modulus of the sequence (s_n) of partial sums, $s_n := \sum_{i=0}^n a_i$.

PROPOSITION 4.3. *Let (X, d, W) be a UCW-hyperbolic space, $C \subseteq X$ be a convex subset and $T : C \rightarrow C$ nonexpansive with $\text{Fix}(T) \neq \emptyset$. Assume that $\sum_{n=0}^{\infty} \lambda_n(1 - \lambda_n)$ is divergent. Then $\liminf_n d(x_n, Ty_n) = 0$ for all $x \in C$.*

Furthermore, if η is a monotone modulus of uniform convexity and $\theta : \mathbb{N} \rightarrow \mathbb{N}$ is a rate of divergence for $\sum_{n=0}^{\infty} \lambda_n(1 - \lambda_n)$, then

for all $x \in C, \varepsilon > 0, k \in \mathbb{N}$ there exists $N \in \mathbb{N}$ such that

$$(12) \quad k \leq N \leq h(\varepsilon, k, \eta, b, \theta) \text{ and } d(x_N, Ty_N) < \varepsilon,$$

where

$$h(\varepsilon, k, \eta, b, \theta) := \begin{cases} \theta\left(\left\lceil \frac{b+1}{\varepsilon \cdot \eta\left(b, \frac{\varepsilon}{b}\right)} \right\rceil + k\right) & \text{for } \varepsilon \leq 2b, \\ k & \text{otherwise,} \end{cases}$$

with $b > 0$ such that $b \geq d(x, p)$ for some $p \in \text{Fix}(T)$.

PROOF. Let $x \in C, p \in \text{Fix}(T)$ and $b > 0$ be such that $d(x, p) \leq b$. Since $(d(x_n, p))$ is decreasing, it follows that $d(x_n, p) \leq d(x, p) \leq b$ for all $n \in \mathbb{N}$.

Let $\varepsilon > 0, k \in \mathbb{N}$ and $\theta : \mathbb{N} \rightarrow \mathbb{N}$ be as in the hypothesis. We shall prove the existence of N satisfying (12), which implies $\liminf_n d(x_n, Ty_n) = 0$.

First, let us remark that $d(x_n, Ty_n) \leq 2d(x_n, p) \leq 2b$ for all $n \in \mathbb{N}$, hence the case $\varepsilon > 2b$ is obvious. Let us consider $\varepsilon < 2b$ and denote

$$P := \left\lceil \frac{b+1}{\varepsilon \eta\left(b, \frac{\varepsilon}{b}\right)} \right\rceil,$$

so $h(\varepsilon, k, \eta, b, \theta) := \theta(P + k) \geq P + k > k$.

Assume by contradiction that $d(x_n, Ty_n) \geq \varepsilon$ for all $n = k, \dots, \theta(P+k)$. Since $b \geq d(x_n, p) \geq \frac{d(x_n, Ty_n)}{2} \geq \frac{\varepsilon}{2}$, we can apply Lemma 4.2.(i) with $\beta := \tilde{\beta} := b, \gamma := \frac{\varepsilon}{2}$ and $a := \varepsilon$ to obtain that for all $n = k, \dots, \theta(P+k)$,

$$(13) \quad d(x_{n+1}, p) \leq d(x_n, p) - \varepsilon \lambda_n (1 - \lambda_n) \eta \left(b, \frac{\varepsilon}{b} \right).$$

Adding (13) for $n = k, \dots, \theta(P+k)$, it follows that

$$\begin{aligned} d(x_{\theta(P+k)+1}, p) &\leq d(x_k, p) - \varepsilon \eta \left(b, \frac{\varepsilon}{b} \right) \sum_{n=k}^{\theta(P+k)} \lambda_n (1 - \lambda_n) \\ &\leq b - \varepsilon \eta \left(b, \frac{\varepsilon}{b} \right) \cdot P \leq b - (b+1) = -1, \end{aligned}$$

that is, a contradiction. We have used the fact that

$$\begin{aligned} \sum_{n=k}^{\theta(P+k)} \lambda_n (1 - \lambda_n) &= \sum_{n=0}^{\theta(P+k)} \lambda_n (1 - \lambda_n) - \sum_{n=0}^{k-1} \lambda_n (1 - \lambda_n) \\ &\geq \sum_{n=0}^{\theta(P+k)} \lambda_n (1 - \lambda_n) - k \geq (P+k) - k = P. \end{aligned}$$

□

As an immediate consequence of the above proposition, we get a rate of asymptotic regularity for the Krasnoselski-Mann iterates, similar with the one obtained in [20, Theorem 1.4].

COROLLARY 4.4. *Let $(X, d, W), \eta, C, T, b, (\lambda_n), \theta$ be as in the hypotheses of Proposition 4.3 and assume that (x_n) is the Krasnoselski-Mann iteration starting with x , defined by (5).*

Then $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ for all $x \in C$ and

$$(14) \quad \forall \varepsilon > 0 \forall n \geq \Phi(\varepsilon, \eta, b, \theta) (d(x_n, Tx_n) < \varepsilon),$$

where $\Phi(\varepsilon, \eta, b, \theta) := h(\varepsilon, 0, \eta, b, \theta)$, with h defined as above.

PROOF. Applying Proposition 4.3 with $s_n := 0$ (hence $y_n = x_n$) and $k := 0$, we get the existence of $N \leq \Phi(\varepsilon, \eta, b, \theta)$ such that $d(x_N, Tx_N) < \varepsilon$. Use the fact that $(d(x_n, Tx_n))$ is decreasing to get (14). □

PROPOSITION 4.5. *In the hypotheses of the above proposition, assume moreover that $\limsup_n s_n < 1$. Then $\liminf_n d(x_n, Tx_n) = 0$ for all $x \in C$.*

Furthermore, if $L, N_0 \in \mathbb{N}$ are such that $s_n \leq 1 - \frac{1}{L}$ for all $n \geq N_0$, then for all $x \in C, \varepsilon > 0, k \in \mathbb{N}$ there exists $N \in \mathbb{N}$ satisfying

$$(15) \quad k \leq N \leq \Psi(\varepsilon, k, \eta, b, \theta, L, N_0) \text{ and } d(x_N, Tx_N) < \varepsilon,$$

where $\Psi(\varepsilon, k, \eta, b, \theta, L, N_0) := h\left(\frac{\varepsilon}{L}, k + N_0, \eta, b, \theta\right)$, h defined as in Proposition 4.3.

PROOF. Let $x \in C, \varepsilon > 0, k \in \mathbb{N}$. Applying Proposition 4.3 for $k + N_0$ and $\frac{\varepsilon}{L}$, we get the existence of N such that $N_0 \leq k + N_0 \leq N \leq h\left(\frac{\varepsilon}{L}, k + N_0, \eta, b, \theta\right) = \Psi(\varepsilon, k, \eta, b, \theta, L, N_0)$ and $d(x_N, Ty_N) < \frac{\varepsilon}{L}$. Using (9), the hypothesis gives

$$d(x_N, Tx_N) \leq \frac{1}{1 - s_N} d(x_N, Ty_N) < \frac{L\varepsilon}{L} = \varepsilon.$$

□

As a corollary, we obtain an approximate fixed point bound for the nonexpansive mapping T .

COROLLARY 4.6. *In the hypotheses of Proposition 4.5,*

$$(16) \quad \forall \varepsilon > 0 \exists N \leq \Phi(\varepsilon, \eta, b, \theta, L, N_0) \left(d(x_N, Tx_N) < \varepsilon \right),$$

where $\Phi(\varepsilon, \eta, b, \theta, L, N_0) := \Psi(\varepsilon, 0, \eta, b, \theta, L, N_0)$, with Ψ defined as above.

We are ready now to prove the main result of this section.

THEOREM 4.7. *Let C be a convex subset of a UCW-hyperbolic space (X, d, W) and $T : C \rightarrow C$ nonexpansive with $\text{Fix}(T) \neq \emptyset$. Assume that $\sum_{n=0}^{\infty} \lambda_n(1 - \lambda_n)$ diverges, $\limsup_n s_n < 1$ and $\sum_{n=0}^{\infty} s_n(1 - \lambda_n)$ converges. Then $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ for all $x \in C$.*

Furthermore, if η is a monotone modulus of uniform convexity, θ is a rate of divergence for $\sum_{n=0}^{\infty} \lambda_n(1 - \lambda_n)$, L, N_0 are such that $s_n \leq 1 - \frac{1}{L}$ for all $n \geq N_0$ and γ is a Cauchy modulus for $\sum_{n=0}^{\infty} s_n(1 - \lambda_n)$, then for all $x \in C$,

$$(17) \quad \forall \varepsilon > 0 \forall n \geq \Phi(\varepsilon, \eta, b, \theta, L, N_0, \gamma) \left(d(x_n, Tx_n) < \varepsilon \right),$$

where

$$\Phi(\varepsilon, \eta, b, \theta, L, N_0, \gamma) := \begin{cases} \theta \left(\left\lceil \frac{2L(b+1)}{\varepsilon \cdot \eta \left(b, \frac{\varepsilon}{2Lb} \right)} \right\rceil + \gamma \left(\frac{\varepsilon}{8b} \right) + N_0 + 1 \right) & \text{for } \varepsilon \leq 4Lb, \\ \gamma \left(\frac{\varepsilon}{8b} \right) + N_0 + 1 & \text{otherwise,} \end{cases}$$

with $b > 0$ such that $b \geq d(x, p)$ for some $p \in \text{Fix}(T)$.

PROOF. Let $x \in C, p \in \text{Fix}(T)$ and $b > 0$ be such that $d(x, p) \leq b$ and let us denote $\alpha_n := \sum_{i=0}^n s_i(1 - \lambda_i)$. Since $d(x_n, Tx_n) \leq 2d(x_n, p) \leq 2b$ for all $n \in \mathbb{N}$, by (11) we get that for all $n \in \mathbb{N}$,

$$d(x_{n+1}, Tx_{n+1}) \leq (1 + 2s_n(1 - \lambda_n))d(x_n, Tx_n) \leq d(x_n, Tx_n) + 4bs_n(1 - \lambda_n);$$

hence for all $m \in \mathbb{N}, n \geq 1$,

$$d(x_{m+n}, Tx_{m+n}) \leq d(x_n, Tx_n) + 4b(\alpha_{n+m-1} - \alpha_{n-1}).$$

Let $\varepsilon > 0$. Applying Proposition 4.5 with $\frac{\varepsilon}{2}$ and $k := \gamma(\varepsilon/8b) + 1$, we get $N \in \mathbb{N}$ such that $d(x_N, Tx_N) < \frac{\varepsilon}{2}$ and

$$\begin{aligned} \gamma(\varepsilon/8b) + 1 \leq N &\leq \Psi \left(\frac{\varepsilon}{2}, \gamma(\varepsilon/8b) + 1, b, \theta, L, N_0 \right) \\ &= h \left(\frac{\varepsilon}{2L}, \gamma(\varepsilon/8b) + 1 + N_0, \eta, b, \theta \right) \\ &= \Phi(\varepsilon, \eta, b, \theta, L, N_0, \gamma). \end{aligned}$$

Since γ is a Cauchy modulus for (α_n) , it follows that for all $m \in \mathbb{N}$,

$$\alpha_{m+\gamma(\varepsilon/8b)} - \alpha_{\gamma(\varepsilon/8b)} = |\alpha_{m+\gamma(\varepsilon/8b)} - \alpha_{\gamma(\varepsilon/8b)}| < \frac{\varepsilon}{8b}.$$

Let now $n \geq \Phi(\varepsilon, \eta, b, \theta, L, N_0, \gamma) \geq N$, hence $n = N + p = \gamma(\varepsilon/8b) + 1 + q$ for some $p, q \in \mathbb{N}$. It follows that

$$\begin{aligned} d(x_n, Tx_n) &= d(x_{N+p}, Tx_{N+p}) \leq d(x_N, Tx_N) + 4b(\alpha_{N+p-1} - \alpha_{N-1}) \\ &= d(x_N, Tx_N) + 4b(\alpha_{\gamma(\varepsilon/8b)+q} - \alpha_{N-1}) \\ &< \frac{\varepsilon}{2} + 4b(\alpha_{\gamma(\varepsilon/8b)+q} - \alpha_{\gamma(\varepsilon/8b)}) \\ &\quad \text{since } N-1 \geq \gamma(\varepsilon/8b), \text{ so } \alpha_{N-1} \geq \alpha_{\gamma(\varepsilon/8b)} \\ &< \frac{\varepsilon}{2} + 4b \cdot \frac{\varepsilon}{8b} = \varepsilon, \end{aligned}$$

since γ is a Cauchy modulus for (α_n) . \square

REMARK 4.8. *In the hypotheses of Theorem 4.7, assume, moreover, that $\eta(r, \varepsilon)$ can be written as $\eta(r, \varepsilon) = \varepsilon \cdot \tilde{\eta}(r, \varepsilon)$ such that $\tilde{\eta}$ increases with ε (for a fixed r). Then the bound $\Phi(\varepsilon, \eta, b, \theta, L, N_0, \gamma)$ can be replaced for $\varepsilon \leq 4Lb$ with*

$$\tilde{\Phi}(\varepsilon, \eta, b, \theta, L, N_0, \gamma) = \theta \left(\left\lceil \frac{L(b+1)}{\varepsilon \cdot \tilde{\eta}\left(b, \frac{\varepsilon}{2Lb}\right)} \right\rceil + \gamma\left(\frac{\varepsilon}{8b}\right) + N_0 + 1 \right).$$

PROOF. As we have seen in the proof of Theorem 4.7,

$$\Phi(\varepsilon, \eta, b, \theta, L, N_0, \gamma) = h\left(\frac{\varepsilon}{2L}, \gamma\left(\frac{\varepsilon}{8b}\right) + 1 + N_0, \eta, b, \theta\right),$$

where h is defined as in Proposition 4.3. It is easy to see that using the extra assumptions on η , $h(\varepsilon, k, \eta, b, \theta)$ can be replaced for $\varepsilon < 2b$ with

$$\tilde{h}(\varepsilon, k, \eta, b, \theta) := \theta \left(\left\lceil \frac{b+1}{2\varepsilon \cdot \tilde{\eta}\left(b, \frac{\varepsilon}{b}\right)} \right\rceil + k \right).$$

Just define $P := \left\lceil \frac{b+1}{2\varepsilon \cdot \tilde{\eta}\left(b, \frac{\varepsilon}{b}\right)} \right\rceil$ and follow the proof of Proposition 4.3 using Lemma 4.2.(ii) (with $\delta := b, a := \varepsilon$) instead of Lemma 4.2.(i). \square

COROLLARY 4.9. *Let (X, d, W) be a complete UCW-hyperbolic space, $C \subseteq X$ a convex closed bounded subset with diameter d_C and $T : C \rightarrow C$ nonexpansive. Assume that $\eta, (\lambda_n), (s_n), \theta, L, N_0, \gamma$ are as in the hypotheses of Theorem 4.7.*

Then $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ for all $x \in C$ and, moreover,

$$\forall \varepsilon > 0 \forall n \geq \Phi(\varepsilon, \eta, d_C, \theta, L, N_0, \gamma) (d(x_n, Tx_n) < \varepsilon),$$

where $\Phi(\varepsilon, \eta, d_C, \theta, L, N_0, \gamma)$ is defined as in Theorem 4.7 by replacing b with d_C .

PROOF. We can apply Corollary 3.6 to get that $\text{Fix}(T) \neq \emptyset$. Moreover, $d(x, p) \leq d_C$ for any $x \in C, p \in \text{Fix}(T)$, hence we can take $b := d_C$ in Theorem 4.7. \square

Thus, for bounded C , we get an effective rate of asymptotic regularity which depends on the error ε , on the modulus of uniform convexity η , on the diameter d_C of C , and on $(\lambda_n), (s_n)$ via θ, L, N_0, γ , but does not depend on the nonexpansive mapping T , the starting point $x \in C$ of the iteration or other data related with C and X .

The rate of asymptotic regularity can be further simplified in the case of constant $\lambda_n := \lambda \in (0, 1)$.

COROLLARY 4.10. *Let $(X, d, W), \eta, C, d_C, T$ be as in the hypotheses of Corollary 4.9. Assume that $\lambda_n := \lambda \in (0, 1)$ for all $n \in \mathbb{N}$.*

Furthermore, let L, N_0 be such that $s_n \leq 1 - \frac{1}{L}$ for all $n \geq N_0$ and assume that the series $\sum_{n=0}^{\infty} s_n$ converges with Cauchy modulus δ .

Then for all $x \in C$,

$$(18) \quad \forall \varepsilon > 0 \forall n \geq \Phi(\varepsilon, \eta, d_C, \lambda, L, N_0, \delta) \left(d(x_n, Tx_n) < \varepsilon \right),$$

where

$$\Phi(\varepsilon, \eta, d_C, \lambda, L, N_0, \delta) := \begin{cases} \left\lceil \frac{1}{\lambda(1-\lambda)} \cdot \frac{2L(d_C+1)}{\varepsilon \cdot \eta\left(d_C, \frac{\varepsilon}{2Ld_C}\right)} \right\rceil + M & \text{for } \varepsilon \leq 4Ld_C, \\ M & \text{otherwise,} \end{cases}$$

with $M := \delta\left(\frac{\varepsilon}{8d_C(1-\lambda)}\right) + N_0 + 1$.

Moreover, if $\eta(r, \varepsilon)$ can be written as $\eta(r, \varepsilon) = \varepsilon \cdot \tilde{\eta}(r, \varepsilon)$ such that $\tilde{\eta}$ increases with ε (for a fixed r), then the bound $\Phi(\varepsilon, \eta, d_C, \lambda, L, N_0, \delta)$ can be replaced for $\varepsilon \leq 4Ld_C$ with

$$\Phi(\varepsilon, \eta, d_C, \lambda, L, N_0, \delta) = \left\lceil \frac{1}{\lambda(1-\lambda)} \cdot \frac{L(d_C+1)}{\varepsilon \cdot \tilde{\eta}\left(d_C, \frac{\varepsilon}{2Ld_C}\right)} \right\rceil + M.$$

PROOF. It is easy to see that

$$\theta : \mathbb{N} \rightarrow \mathbb{N}, \quad \theta(n) = \left\lceil \frac{n}{\lambda(1-\lambda)} \right\rceil$$

is a rate of divergence for $\sum_{n=0}^{\infty} \lambda(1-\lambda)$. Moreover,

$$\gamma : (0, \infty) \rightarrow \mathbb{N}, \quad \gamma(\varepsilon) = \delta\left(\frac{\varepsilon}{1-\lambda}\right)$$

is a Cauchy modulus for $\sum_{n=0}^{\infty} s_n(1-\lambda)$. Apply now Corollary 4.9 and Remark 4.8. \square

As we have seen in Section 2, CAT(0) spaces are UCW-hyperbolic spaces with a modulus of uniform convexity $\eta(r, \varepsilon) := \frac{\varepsilon^2}{8}$, which has the form required in Remark 4.8. Thus, the above result can be applied to CAT(0) spaces.

COROLLARY 4.11. *Let X be a CAT(0) space, $C \subseteq X$ a bounded convex closed subset with diameter d_C and $T : C \rightarrow C$ nonexpansive. Assume that $\lambda_n = \lambda \in (0, 1)$ for all $n \in \mathbb{N}$ and $L, N_0, (s_n), \delta$ are as in the hypotheses of Corollary 4.10*

Then $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ for all $x \in C$ and, moreover,

$$(19) \quad \forall \varepsilon > 0 \forall n \geq \Phi(\varepsilon, d_C, \lambda, L, N_0, \delta) \left(d(x_n, Tx_n) < \varepsilon \right),$$

where

$$\Phi(\varepsilon, d_C, \lambda, L, N_0, \delta) := \begin{cases} \left\lceil \frac{D}{\varepsilon^2} \right\rceil + M, & \text{for } \varepsilon \leq 4Ld_C, \\ M & \text{otherwise,} \end{cases}$$

with $M := \delta\left(\frac{\varepsilon}{8d_C(1-\lambda)}\right) + N_0 + 1$, $D = \frac{16L^2 d_C(d_C+1)}{\lambda(1-\lambda)}$.

References

- [1] J. Borwein, S. Reich and I. Shafrir, *Krasnoselskii-Mann iterations in normed spaces*, Canad. Math. Bull. **35** (1992), 21–28.
- [2] M. Bridson and A. Haefliger, *Metric spaces of non-positive curvature, Grundlehren der Mathematischen Wissenschaften*, vol. 319, Springer-Verlag, 1999.
- [3] J.A. Clarkson, *Uniformly convex spaces*, Trans. Amer. Math. Soc. **40** (1936), 396–414.
- [4] S. Dhompongsa and B. Panyanak, *On Δ -convergence theorems in $CAT(0)$ spaces*, Computers & Mathematics with Applications **56** (2008), no. 10, 2572–2579.
- [5] W. G. Dotson, Jr. and W. R. Mann, *A generalized corollary of the Browder-Kirk fixed point theorem*, Pacific J. Math. **26** (1968), 455–459.
- [6] M. Edelstein, *The construction of an asymptotic center with a fixed-point property*, Bull. Amer. Math. Soc. **78** (1972), 206–208.
- [7] M. Edelstein, *Fixed point theorems in uniformly convex Banach spaces*, Proc. Amer. Math. Soc. **44** (1974), 369–374.
- [8] K. Goebel and W.A. Kirk, *Iteration processes for nonexpansive mappings*, in: Topological Methods in Nonlinear Functional Analysis (S. P. Singh, S. Thomeier, B. Watson, eds.)(Toronto, 1982), pp. 115–123, Contemp. Math. vol. 21, Amer. Math. Soc., 1983.
- [9] K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Monographs and Textbooks in Pure and Applied Mathematics vol. 83, Marcel Dekker Inc., 1984.
- [10] C. W. Groetsch, *A note on segmenting Mann iterates*, J. Math. Anal. and Appl. **40** (1972), 369–372.
- [11] S. Ishikawa, *Fixed points by a new iteration method*, Proc. Amer. Math. Soc. **44** (1974), 147–150.
- [12] S. Itoh, *Some fixed point theorems in metric spaces*, Fund. Math. **102** (1979), 109–117.
- [13] W.A. Kirk, *Krasnosel'skii iteration process in hyperbolic spaces*, Numer. Funct. Anal. and Optimiz. **4** (1982), 371–381.
- [14] U. Kohlenbach, *A quantitative version of a theorem due to Borwein-Reich-Shafrir*, Numer. Funct. Anal. Optim. **22** (2001), 641–656.
- [15] U. Kohlenbach, *Some logical metatheorems with applications in functional analysis*, Trans. Amer. Math. Soc. **357** (2005), 89–128.
- [16] U. Kohlenbach, *Applied Proof Theory: Proof Interpretations and their Use in Mathematics*, Springer Monographs in Mathematics, Springer-Verlag, 2008.
- [17] U. Kohlenbach and L. Leuştean, *Mann iterates of directionally nonexpansive mappings in hyperbolic spaces*, Abstract and Applied Analysis **2003** (2003), 449–477.
- [18] U. Kohlenbach and L. Leuştean, *Asymptotically nonexpansive mappings in uniformly convex hyperbolic spaces*, arXiv:0707.1626v2 [math.FA], 2007, to appear in J. European Math. Soc.
- [19] M. A. Krasnoselski, *Two remarks on the method of successive approximation*, Usp. Math. Nauk (N.S.) **10** (1955), 123–127 (Russian).
- [20] L. Leuştean, *A quadratic rate of asymptotically regularity for $CAT(0)$ spaces*, J. Math. Anal. Appl. **325** (2007), 386–399.
- [21] L. Leuştean, *Rates of asymptotic regularity for Halpern iterations of nonexpansive mappings*, in Combinatorics and Related Areas. A Collection of Papers in Honor of the 65th Birthday of Ioan Tomescu (C.S. Calude, G. Ştefănescu, and M. Zimand, eds.), Journal of Universal Computer Science **13** (2007), 1680–1691.
- [22] W. R. Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. **4** (1953), 506–510.
- [23] S. Reich, *Fixed point iterations of nonexpansive mappings*, Pacific J. Math. **60** (1975), 195–198.
- [24] S. Reich and I. Shafrir, *Nonexpansive iterations in hyperbolic spaces*, Nonlinear Analysis **15** (1990), 537–558.
- [25] H. Schaefer, *Über die Method sukzessive Approximationen*, J. Deutsch Math. Verein **59** (1957), 131–140.
- [26] W. Takahashi, *A convexity in metric space and nonexpansive mappings I*, Kodai Math. Sem. Rep. **22** (1970), 142–149.
- [27] K.-K. Tan and H.-K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl. **178** (1993), 301–308.

DEPARTMENT OF MATHEMATICS, TECHNISCHE UNIVERSITÄT DARMSTADT, SCHLOSSGARTEN-
STRASSE 7, 64289 DARMSTADT, GERMANY

AND

INSTITUTE OF MATHEMATICS "SIMION STOILOW" OF THE ROMANIAN ACADEMY, P.O. BOX 1-764,
014700 BUCHAREST, ROMANIA

E-mail address: `leustean@mathematik.tu-darmstadt.de`